

Universally measurable sets may all be Δ_2^1

Paul B. Larson*
Miami University
Oxford, Ohio USA

Saharon Shelah†
Hebrew University of Jerusalem
Rutgers University.

March 5, 2024

Abstract

We produce a forcing extension of the constructible universe \mathbf{L} in which every universally measurable set of reals is Δ_2^1 , partially answering question CG from David Fremlin's problem list [5]. The analogous result for category holds in the same model.

We let ω^ω denote the Baire space, the set of functions from ω to ω , and refer to its elements as *reals*. A subset of a Polish space X is said to be *universally measurable* if it is measured by the completion of any σ -additive Borel measure on X . Equivalently, $A \subseteq X$ is universally measurable if and only if $f^{-1}[A]$ is Lebesgue measurable whenever $f: \omega^\omega \rightarrow X$ is a Borel function (see [7, 11], and 434D of [4], for instance). This characterization induces the corresponding notion for category: we will say that a set $A \subseteq X$ is *universally categorical* if and only if $f^{-1}[A]$ has the property of Baire whenever $f: \omega^\omega \rightarrow X$ is a Borel function. (The term *universally Baire* has already been established with a different meaning [2], implying both universal measurability and universal categoricity.) The collections of the universally measurable subsets of X and the universally categorical subsets of X are both σ -algebras on X .

A subset A of a Polish space X is Δ_2^1 if A and $X \setminus A$ are continuous images of coanalytic sets. We refer the reader to [7] for background on this definition and for information on the projective sets in general. In this paper we show how to force over any model of the form $\mathbf{L}[a]$, where $a \subseteq \omega$, to produce a model in which all subsets of ω^ω which are either universally measurable or universally categorical are Δ_2^1 . Since all uncountable Polish spaces are Borel-isomorphic (see Theorem 17.41 of [7]), this implies the same fact for subsets of any Polish

*The research of the first author was supported in part by NSF Grants DMS-1201494 and DMS-1764320.

†The research of the second author was partially supported by NSF grant number 1101597, and by European Research Council grant 338821. This is paper number 1178 on the second author's list. The authors thank the referee for closely reading this paper and making many helpful suggestions.

space. The following then is the main theorem of this paper (Theorem 5.1 gives a more explicit statement).

{specthrm}

Theorem 0.1. *If, for some $a \subseteq \omega$, $\mathbf{V}=\mathbf{L}[a]$, then there is a proper forcing extension in which every universally measurable subset of any Polish space is Δ_2^1 , and every universally categorical subset of any Polish space is Δ_2^1 .*

The result for universally measurable sets answers part of problem CG on David Fremlin's problem list [5].

Since there are only continuum many Δ_2^1 sets, our result also strengthens (modulo the anti-large cardinal hypothesis $\mathbf{V}=\mathbf{L}[a]$) a previous result of the authors with Itay Neeman [9], which showed the consistency of the statement that the set of universally measurable sets has the same cardinality as ω^ω . We note that (unlike the results in [9]) some anti-large cardinal hypothesis is needed for the results in this paper, since the existence of infinitely many Woodin cardinals for instance implies that every projective set of reals is universally measurable [17], and there are (assuming ZF) projective sets which are not Δ_2^1 (see Theorem 37.7 of [7]).

1 Outline of the proof

{outlinesec}

The proof of Theorem 0.1 is an application of forcing machinery developed by the second author and his collaborators (especially [14, 15], but we also make use of results from [8]). The proof proceeds by forcing over a model of the form $\mathbf{L}[a]$ (for any $a \subseteq \omega$) with a countable support iteration of proper partial orders, and makes use of the following theorem, which is Theorem III.4.1 in [16].

{shthrm}

Theorem 1.1. *Suppose that κ is a regular cardinal such that $\mu^{\aleph_0} < \kappa$ for all $\mu < \kappa$ and that $\bar{\mathbb{P}} = \langle \mathbb{P}_\alpha, \dot{Q}_\beta : \alpha \leq \kappa, \beta < \kappa \rangle$ is a countable support iteration such that each \mathbb{P}_α forces the corresponding \dot{Q}_α to be a proper forcing of cardinality less than κ . Then \mathbb{P}_κ is κ -c.c., and for each $\alpha < \kappa$, \mathbb{P}_α has a dense subset of cardinality less than κ . Furthermore, for all $\alpha < \kappa$, \mathbb{P}_α forces that $2^{\aleph_0} < \kappa$.*

We will apply this theorem with ground models of the form $\mathbf{L}[a]$ for $a \subseteq \omega$, which satisfy the Generalized Continuum Hypothesis. Although we could let κ be any regular cardinal with $\kappa^{<\kappa} = \kappa$ we will restrict ourselves to the case $\kappa = \omega_2^{\mathbf{V}}$. Each step of our iterations will be an (ω, ∞) -distributive partial order of cardinality continuum (see Definition 2.2), and will force the Continuum Hypothesis (CH) to hold. To see that Theorem 1.1 applies, we need to know that each \mathbb{P}_α preserves the statement that $2^{\aleph_0} < \kappa$. This is not hard to show directly for the partial orders we consider, but it also follows by applying Theorem 1.1 to the modified iteration where each \dot{Q}_α is the \mathbb{P}_α -name for either our original \dot{Q}_α if CH holds, and the trivial forcing if it fails. The theorem then implies that the second case never holds. We then have from Theorem 1.1 that each \mathbb{P}_α (in our original, intended iteration) will have a dense subset of cardinality less than κ , and will therefore preserve the statement $2^{\aleph_1} \leq \kappa$.

We will use the following definition.

1.2 Definition. Given a partial order P and a σ -algebra \mathcal{A} on ω^ω we say that P is \mathcal{A} -representing if for each $A \in \mathcal{A}$ (in \mathbf{V}), every element of ω^ω in any forcing extension by P is in a Borel set with a code in \mathbf{V} which is either contained in or disjoint from A .

Each tail of the iterations we consider here is \mathcal{A} -representing when \mathcal{A} is either the algebra of universally measurable or universally categorical sets. The overall strategy of our proof is then summarized by the following proposition.

Proposition 1.3. *Suppose that $\bar{\mathbb{P}}$ is a forcing iteration of length κ satisfying the conditions in Theorem 1.1. Let $G \subseteq \mathbb{P}_\kappa$ be a \mathbf{V} -generic filter, and for each $\alpha < \kappa$ let G_α be the restriction of G to \mathbb{P}_α . Let \mathcal{A}_α ($\alpha \leq \kappa$) be such that either each \mathcal{A}_α is the set of universally measurable subsets of ω^ω in $\mathbf{V}[G_\alpha]$ or each \mathcal{A}_α is the set of universally categorical subsets of ω^ω in $\mathbf{V}[G_\alpha]$. Suppose that each tail $\mathbb{P}_{[\alpha, \kappa]}$ of $\bar{\mathbb{P}}$ is \mathcal{A}_α -representing. Then for each $A \in \mathcal{A}_\kappa$ there exists an $\alpha < \kappa$ such that A is equal to the union of all the Borel subsets of A in $\mathbf{V}[G]$ having codes in $\mathbf{V}[G_\alpha]$.*

Proof. Fixing A , we have that A is the realization of a \mathbb{P}_κ -name \dot{A} in the ground model \mathbf{V} . We have the following facts:

- each \mathbb{P}_α preserves the statement $2^{\aleph_0} < \kappa$;
- \mathbb{P}_κ preserves the regularity of κ ;
- for each $A \subseteq \omega^\omega$ the statement that A is universally measurable and the statement that A is universally categorical are both \prod_2^1 in A .

Together these imply that for club many $\alpha < \kappa$ (more importantly, at least one), the following hold.

- For each \mathbb{P}_α -name σ for an element of ω^ω , some element of G_α decides the statement $\sigma \in \dot{A}$ (with respect to \mathbb{P}_κ , with σ interpreted as a \mathbb{P}_κ -name).
- For each Borel set B in $V[G]$ with a code in $\mathbf{V}[G_\alpha]$, if $B \cap \dot{A}_G$ is nonempty, then so is $B \cap \dot{A}_G \cap \mathbf{V}[G_\alpha]$.
- For each Borel set B in $V[G]$ with a code in $\mathbf{V}[G_\alpha]$, if $B \setminus \dot{A}_G$ is nonempty, then so is $(B \setminus \dot{A}_G) \cap \mathbf{V}[G_\alpha]$.
- If \dot{A}_G is universally measurable in $\mathbf{V}[G]$, then $\dot{A}_G \cap \mathbf{V}[G_\alpha]$ is universally measurable in $\mathbf{V}[G_\alpha]$.
- If \dot{A}_G is universally categorical in $\mathbf{V}[G]$, then $\dot{A}_G \cap \mathbf{V}[G_\alpha]$ is universally categorical in $\mathbf{V}[G_\alpha]$.

To finish the proof, let x be an element of A . Since $A \cap \mathbf{V}[G_\alpha]$ is in \mathcal{A}_α , and the tail $\mathbb{P}_{[\alpha, \kappa]}$ of $\bar{\mathbb{P}}$ is \mathcal{A}_α -representing, x is a member of a Borel set B with a code in $\mathbf{V}[G_\alpha]$ such that $B \cap \mathbf{V}[G_\alpha]$ is either contained in or disjoint from A . Since B is not disjoint from A , $B \cap \mathbf{V}[G_\alpha]$ is not either. So $B \cap \mathbf{V}[G_\alpha]$ is contained in A , and B is also. \square

The forcing construction in this paper produces a model in which every set of reals of cardinality \aleph_1 is Σ_2^1 (the fact that each \mathbb{P}_α preserves the inequality $2^{\aleph_1} \leq \kappa$ makes this possible with an iteration of length κ). Given this, any $A \subseteq \omega^\omega$ with the property that A and $\omega^\omega \setminus A$ are both unions of \aleph_1 -many Borel sets is Δ_2^1 (see Remark 2.5).

{reducerem}

1.4 Remark. The outline in this section reduces the proof of the main theorem to establishing the following regarding the iterations $\langle \mathbb{P}_\alpha, \dot{Q}_\beta : \alpha \leq \kappa, \beta < \kappa \rangle$ considered in this paper (iterations as in Definition 3.1 and their tails):

- each \mathbb{P}_α forces that \dot{Q}_α is a proper (ω, ∞) -distributive forcing of cardinality at most 2^{\aleph_0} making CH hold (established in Remark 2.3 and Lemma 2.4);
- \mathbb{P}_κ forces that every subset of ω^ω of cardinality \aleph_1 is Σ_2^1 (shown in Lemma 2.6);
- each tail of the iteration \mathbb{P}_κ is \mathcal{A} -representing when \mathcal{A} is either the set of universally measurable subsets of ω^ω or the set of universally categorical subsets of ω^ω . This is shown in Lemma 5.2.

At the end of the paper we prove one additional result not directly related to the main theorem. The paper [9] introduced the following notion: given a ground model set $A \subseteq \omega^\omega$, the *Borel reinterpretation* of A in a forcing extension is the union of all the ground model Borel sets contained in A , each reinterpreted in the extension. This offered a characterization of the universally measurable sets as the sets $A \subseteq \omega^\omega$ with the property that the Borel reinterpretations of A and $\omega^\omega \setminus A$ are complements in any extension by random forcing (in particular, random forcing is \mathcal{A} -representing, when \mathcal{A} is the collection of universally measurable sets). Similarly, the universally categorical sets are the sets $A \subseteq \omega^\omega$ with the property that the Borel reinterpretations of A and $\omega^\omega \setminus A$ are complements in any extension by Cohen forcing.

Given a definable σ -ideal \mathcal{A} on ω^ω , we say that a partial order P has the *\mathcal{A} -reinterpretation property* if it is \mathcal{A} -representing, and, in addition the Borel reinterpretations of members of \mathcal{A} in the ground model are in \mathcal{A} as defined in forcing extensions by P . Many forcings have this property for the universal measurable sets, including random forcing and Sacks forcing; see [10]. Theorem 6.1 shows that the iterations considered in this paper have the \mathcal{A} -reinterpretation property when \mathcal{A} is the set of universally measurable subsets of ω^ω . This is in some sense a negative result: an iteration forcing the statement “every universally measurable set has the property of Baire” (whose consistency is still an open question) cannot have the reinterpretation property for universally measurable sets if, for instance, it is applied to a model of Martin’s Axiom (more generally, to a model with a medial limit).

2 Coding subsets of ω_1 by reals

{stepsec}

We let Ω denote the set of countable limit ordinals. A *ladder system* on ω_1 is a sequence $\langle C_\eta : \eta \in \Omega \rangle$ such that each C_η is a cofinal subset of η of ordertype ω .

For C an infinite set of ordinals and $n \in \omega$, we write $C(n)$ for the unique $\alpha \in C$ such that $|C \cap \alpha| = n$.

Given two sequences s, t , we write $s \sqsubseteq t$ mean that s is an initial segment of t . Given $s \in 2^{<\omega}$, we let $[s]$ denote $\{x \in {}^\omega 2 : s \sqsubseteq x\}$.

{denfundef}

2.1 Definition. We define a *dense function* to be a partial function $F: 2^\omega \rightarrow 2$ such that for each $s \in 2^{<\omega}$, $F[[s] \cap \text{dom}(F)] = 2$.

{podef}

2.2 Definition. We define the forcing $Q_{\bar{C}, F, g}$, where

- $\bar{C} = \langle C_\eta : \eta \in \Omega \rangle$ is a ladder system on ω_1 ;
- $F: 2^\omega \rightarrow 2$ is a dense partial function;
- g is a function from Ω to 2.

The conditions of $Q_{\bar{C}, F, g}$ are the functions p such that,

- the domain of p is a countable ordinal δ_p ;
- the range of p is a subset of 2;
- for all $\eta \in (\delta_p + 1) \cap \Omega$, $\langle p(C_\eta(i)) : i < \omega \rangle \in \text{dom}(F)$ and

$$F(\langle p(C_\eta(i)) : i < \omega \rangle) = g(\eta).$$

The order on $Q_{\bar{C}, F, g}$ is extension.

{popropsrem}

2.3 Remark. Each partial order of the form $Q_{\bar{C}, F, g}$ has cardinality 2^{\aleph_0} , as does its transitive closure (so $Q_{\bar{C}, F, g}$ is in $H(\mathfrak{c}^+)$). Each such partial order also forces $2^{\aleph_0} = \aleph_1$. To see this, note first that by Lemma 2.4 below, $Q_{\bar{C}, F, g}$ adds no new elements of ω^ω . To see that forcing with $Q_{\bar{C}, F, g}$ wellorders $(2^\omega)^{\mathbf{V}}$ in ordertype ω_1 , let \bar{D} be a ladder system on ω_1 such that

- $D_\eta \cap C_\eta = \emptyset$ for each $\eta \in \Omega$ of ordertype greater than ω and
- $\sup\{D_\eta(0) : \eta \in \Omega\} = \omega_1$.

Let $G: \omega_1 \rightarrow 2$ be a V -generic function for $Q_{\bar{C}, F, g}$, and note that each element of $(2^\omega)^{\mathbf{V}}$ is equal to $\langle G(D_\eta(i)) : i < \omega \rangle$ for some $\eta \in \Omega$ (this follows from a standard genericity argument, and we leave the details to the reader).

The following lemma establishes basic properties of the partial orders $Q_{\bar{C}, F, g}$.

{properlem}

Lemma 2.4. *If \bar{C} is a ladder system on ω_1 , $F: 2^\omega \rightarrow 2$ is a partial dense function and g is a function from Ω to 2 then the following hold.*

1. *For each condition $p \in Q_{\bar{C}, F, g}$, each $\gamma < \omega_1$ and each finite partial function s from $\gamma \setminus \delta_p$ to 2, there is a condition $q \leq p$ with $\delta_q \geq \gamma$ and $q \upharpoonright \text{dom}(s) = s$.*
2. *The partial order $Q_{\bar{C}, F, g}$ is proper and (ω, ∞) -distributive.*

Proof. We prove the first part by induction on γ . There is nothing to do in the case where $\gamma \leq \delta_p$. The case where γ is the successor of some ordinal η follows from the induction hypothesis for η along with an extension to add η to the domain of the condition q , agreeing with s if necessary. For the case where γ is a limit ordinal, an application of the induction hypothesis can be used to find an extension p' of p containing s . Since F is dense there is a $y \in 2^\omega$ extending

$$\langle p'(C_\gamma(i)) : i < \omega, C_\gamma(i) < \delta_{p'} \rangle$$

with $F(y) = g(\gamma)$. Applying the induction hypothesis ω any times (with the values $C_\gamma(i) + 1$ for which $C_\gamma(i) \notin \text{dom}(p')$ successively in the role of γ , and the restriction of y to the values $C_\gamma(i)$ in the role of s) gives a condition q as desired.

The argument for the second part is similar. Let p_0 be a condition in $Q_{\bar{C}, F, g}$. Let X be a countable elementary substructure of $H(\aleph_3^+)$ with p_0, \bar{C}, F and g in X . Let $\gamma = X \cap \omega_1$ and let y be an element of 2^ω extending

$$\langle p_0(C_\gamma(i)) : i < \omega, C_\gamma(i) < \delta_{p_0} \rangle$$

with $F(y) = g(\gamma)$. Let R be the set of $q \leq p_0$ such that

$$\langle q(C_\gamma(i)) : i < \omega, C_\gamma(i) < \delta_q \rangle$$

is an initial segment of y .

We claim that for each dense subset D of $Q_{\bar{C}, F, g}$ in X and each $p \in R \cap X$, there is a $q \leq p$ in $R \cap D \cap X$. To see that the claim holds, fix D and p and let $Y \in X$ be a countable elementary substructure of $H((\aleph_2)^+)$ such that $D, p \in Y$. Extend p to a condition $p' \in R \cap Y$ with $C_\gamma \cap Y \subseteq \delta_{p'}$. Then let $q \leq p'$ be an element of $Y \cap D$. It follows from the claim that there exists a condition below p_0 which is in each dense open subset of $Q_{\bar{C}, F, g}$ in X . \square

{aleph1Borelrem}

2.5 Remark. In the forcing extension we produce, each subset of ω^ω which is either universally measurable or universally categorical will be a union of \aleph_1 many Borel sets. Lemma 2.6 below will be used to show that in addition, in this extension, every subset of $\mathcal{P}(\omega)$ of cardinality \aleph_1 is Σ_2^1 . Analytic subsets of ω^ω are naturally coded by elements of $\mathcal{P}(\omega)$ (coding trees; see for instance Section 27A of [7]) in such a way that the set of pairs (x, y) such that $x \subseteq \omega$, $y \in \omega^\omega$ and x is in the set coded by y is analytic. From this one gets that, in our extension, all subsets of ω^ω which are either universally measurable or universally categorical are Σ_2^1 . Since the set of universally measurable sets and the set of universally categorical sets are both closed under complements, it follows that, in this extension, each such set is Δ_2^1 .

Our coding of elements of $[\mathcal{P}(\omega)]^{\aleph_1}$ uses certain iterations of length ω of partial orders of the form $Q_{\bar{C}, F, g}$, which we now define. Let $\pi: \omega \times \omega \rightarrow \omega$ be a fixed recursive bijection, and let π_0 and π_1 be functions from ω to ω such that $\pi(i) = (\pi_0(i), \pi_1(i))$ for all $i < \omega$. We choose π so that $\pi_0(i) < i$ for all $i > 0$. We define, for each triple (\bar{C}, F, g) such that

- \bar{C} is a ladder system on ω_1 ,
- F is a dense function from 2^ω to 2 and
- g is a function from Ω to 2

the following objects recursively on $i < \omega$ (and suppress discussion of the meaning of “canonical name”, trusting the reader to supply her or his preferred definition). Let

- P_0 be the trivial partial order;
- \dot{g}_0 be the canonical P_0 -name for g ;
- \dot{Q}_0 be the canonical P_0 -name for $Q_{\bar{C}, F, g}$;
- for all $i < \omega$,
 - P_{i+1} be $P_i * \dot{Q}_i$;
 - \dot{h}_i be the canonical P_{i+1} -name for the \dot{Q}_i -generic function from ω_1 to 2;
- for all positive $i < \omega$,
 - \dot{g}_i be the canonical P_i -name for the set of pairs (α, k) such that $\alpha \in \Omega$ and $k = \dot{h}_{\pi_0(i), G_i}(\alpha + \pi_1(i))$, where G_i denotes the generic filter for P_i ;
 - \dot{Q}_i be the canonical P_i -name for $Q_{\bar{C}, F, \dot{g}_i}$.

We then let $Q_{\bar{C}, F, g}^*$ denote the full (i.e., countable) support limit of the forcing iteration $\langle P_i, \dot{Q}_i : i < \omega \rangle$.

The purpose of this definition is given in Lemma 2.6 below, but the key point is that, for any $\alpha \in \Omega$, the values $g_i(\alpha)$ ($i < \omega$) determined via π the values $h_i(\alpha + n)$ ($i, n \in \omega$), and similarly, for any $\beta \in \Omega$ and the values of $h_i(\alpha)$ ($i \in \omega, \alpha < \beta$) determine the values $g_i(\beta)$ ($i < \omega$) via F . So relative to the ground model objects F and π , the sequence $\langle h_i \upharpoonright \omega : i < \omega \rangle$ determines the sequence $\langle (g_i, h_i) : i < \omega \rangle$.

Given $a \subseteq \omega$, we define the *canonical ladder system relative to a* to be the set $\bar{C}_a = \{C_\alpha^a : \alpha < \Omega \cap \omega_1^{\mathbf{L}[a]}\}$, where each C_α^a is the constructibly least (in $\mathbf{L}[a]$, relative to a) cofinal subset of the corresponding α of ordertype ω . This defines a ladder system in $\mathbf{L}[a]$ which is a ladder system in \mathbf{V} if and only if $\omega_1^{\mathbf{L}[a]} = \omega_1$. Let Ω' be the set of countable limits of limit ordinals.

{codingitem}

Lemma 2.6. *Let a be a subset of ω such that $\omega_1^{\mathbf{L}[a]} = \omega_1$, and let $F: 2^\omega \rightarrow 2$ be a dense function whose graph is Σ_2^1 in $\mathbf{L}[a]$. Let B be a subset of $\mathcal{P}(\omega)$ of cardinality at most \aleph_1 . There exists a function $g: \Omega \rightarrow 2$ such that, whenever G is a V -generic filter for $Q_{\bar{C}_a, F, g}^*$ and M is an outer model of $\mathbf{V}[G]$, then B is Σ_2^1 in M .*

Proof. It suffices to consider the case where B is nonempty. We fix (and suppress) a Σ_2^1 definition for F , and note that, by Shoenfield absoluteness, in any outer model this definition will define a partial function from ω^ω to 2 containing F .

Let $g: \Omega \rightarrow 2$ be such that

$$B = \{\{k \in \omega : g(\beta + \omega \cdot k) = 0\} : \beta \in \Omega'\},$$

let G be \mathbf{V} -generic for $Q_{\bar{C}_\alpha, F, g}^*$ and let M be an outer model of $\mathbf{V}[G]$. For each $i < \omega$, let h_i be $\dot{h}_{i, G \upharpoonright P_{i+1}}$, where \dot{h}_i is as in the definition of $Q_{\bar{C}, F, g}^*$. We show that, in M , B is Σ_2^1 in a and $\langle h_i \upharpoonright \omega : i < \omega \rangle$. In particular, B is the set of $x \subseteq \omega$ such that there exist

- an element y of ω^ω coding a model of the form $\mathbf{L}_\alpha[a]$, for some countable ordinal α (the wellfoundedness of this model being a Π_1^1 condition on y), such that $\omega_1^{\mathbf{L}_\alpha[a]}$ exists (i.e., some element of α is uncountable in $\mathbf{L}_\alpha[a]$),
- $\beta \in \Omega' \cap \omega_1^{\mathbf{L}_\alpha[a]}$,
- functions $h_i^*: \beta + \omega \cdot \omega \rightarrow 2$ ($i < \omega$) such that, for each $i < \omega$, $h_i^* \upharpoonright \omega = h_i \upharpoonright \omega$ and
- functions $g_i^*: \Omega \cap (\beta + \omega \cdot \omega) \rightarrow 2$ ($i < \omega$)

such that, letting F^* be the function computed in $\mathbf{L}_\alpha[a]$ using our Σ_2^1 definition for F ,

{liftitem}

1. for each $i < \omega$ and each $\gamma \in \Omega \cap (\beta + \omega \cdot \omega)$,
 - $\langle h_i^*(C_\gamma^a(j)) : j < \omega \rangle$ is in the domain of F^* ;
 - $g_i^*(\gamma) = F^*(\langle h_i^*(C_\gamma^a(j)) : j < \omega \rangle)$;
 - if $i > 0$ then $g_i^*(\gamma) = h_{\pi_0(i)}^*(\gamma + \pi_1(i))$;
2. $x = \{k \in \omega : g_0^*(\beta + \omega \cdot k) = 0\}$.

That there exist such objects for each element of B follows from the choice of g and the definition of $Q_{\bar{C}_\alpha, F, g}^*$. Given such objects, the choice of the names \dot{g}_i (and item (1) above) then implies (via an inductive proof on $\gamma \in \Omega \cap (\beta + \omega \cdot \omega)$) that for each $i < \omega$, $g_i^* = \dot{g}_{i, G \upharpoonright P_i} \upharpoonright (\beta + \omega \cdot \omega)$ and $h_i^* = h_i \upharpoonright (\beta + \omega \cdot \omega)$, which, again by the choice of g , implies that the corresponding set x is in B . \square

3 Sequences and trees

{seqtreesecc}

Remarks 1.4 and 2.3 and Lemma 2.6 reduce the proof of Theorem 0.1 to showing that the forcing iterations we consider are \mathcal{A} -representing, where \mathcal{A} is either the collection of universally measurable sets or the collection of universally categorical sets. These iterations will be countable support iterations where each

successor step is a partial order of the form $Q_{\bar{C},F,\dot{g}}$, for a fixed ladder system \bar{C} on ω_1 and a fixed dense function F (moreover, our iterations will consist of segments of the form $Q_{\bar{C},F,\dot{g}}^*$). An additional requirement on F (total pathology) will be introduced in Section 4. We fix the following notation.

{QCFdef}

3.1 Definition. Let \bar{C} be a ladder system on ω_1 and let $F: 2^\omega \rightarrow 2$ be a dense partial function. Let $\mathbf{Q}_{\bar{C},F}$ be the class of sequences of the form

$$\langle \mathbb{P}_\alpha, \dot{Q}_\beta, \dot{g}_\beta, \dot{h}_\beta : \alpha \leq \alpha_*, \beta < \alpha_* \rangle,$$

where

- α_* is an ordinal,
- $\langle \mathbb{P}_\alpha, \dot{Q}_\beta : \alpha \leq \alpha_*, \beta < \alpha_* \rangle$ is a countable support iteration,
- each \dot{g}_β is a P_β -name for a function from Ω to 2,
- each \dot{Q}_β is a P_β -name for the partial order $Q_{\bar{C},F,\dot{g}_\beta}$,
- each \dot{h}_β is a $P_{\beta+1}$ -name for the \dot{Q}_β -generic function.

We say that $\langle \mathbb{P}_\alpha, \dot{Q}_\beta, \dot{g}_\beta, \dot{h}_\beta : \alpha \leq \alpha_*, \beta < \alpha_* \rangle \in \mathbf{Q}_{\bar{C},F}$ is *fully bookkeeping* if for each $\alpha < \alpha_*$ and each \mathbb{P}_α -name \dot{B} for a nonempty set of reals of cardinality at most \aleph_1 , there is a $\alpha' \in [\alpha, \alpha_*)$ such that $1_{\mathbb{P}_{\alpha'}}$ forces that the iteration induced by $\langle \dot{Q}_{\alpha'+n} : n \in \omega \rangle$ will be a forcing of the form $Q_{\bar{C},F,g}^*$ (as computed in the $\mathbb{P}_{\alpha'}$ -extension), with g chosen as in Lemma 2.6 so that $Q_{\bar{C},F,g}^*$ forces B to be Σ_2^1 in its extension and in the \mathbb{P}_{α_*} extension.

In Section 5 we will be building a suitable X -generic condition for some \mathbb{P}_{α_*} as above, where X is a countable elementary submodel of a large enough initial segment of the universe containing our iteration. The first definition below lists useful data that comes with such a situation. The second definition describes a tree of conditions which will guide us to finding our desired condition.

{sdsdef}

3.2 Definition. A *suitable data sequence* is sequence

$$\langle \bar{C}, F, \mathbf{q}, p, X, \bar{Y}, \bar{1}, \bar{D} \rangle$$

such that

- $\bar{C} = \langle C_\alpha : \alpha \in \Omega \rangle$ is a ladder system on ω_1 ;
- $F: 2^\omega \rightarrow 2$ is a dense function;
- $\mathbf{q} = \langle \mathbb{P}_\alpha, \dot{Q}_\beta, \dot{g}_\beta, \dot{h}_\beta : \alpha \leq \alpha_*, \beta < \alpha_* \rangle$ is in $\mathbf{Q}_{\bar{C},F}$;
- p is in \mathbb{P}_{α_*} ;
- $X \prec (H((2^{2^{\aleph_1}})^+), \in)$ is countable, with $\{\bar{C}, F, \mathbf{q}, p\} \in X$;

- $\bar{Y} = \langle Y_k : k \in \omega \rangle$ is an \in -chain of countable elementary substructures of $H((2^{\aleph_1})^+)$, such that
 - $\{\bar{C}, F, \mathbf{q}, p\} \in Y_0$,
 - $X \cap H((2^{\aleph_1})^+) = \bigcup_{k \in \omega} Y_k$,
 - for all $k \in \omega$, $C_{X \cap \omega_1} \cap (Y_{k+1} \setminus Y_k) \neq \emptyset$;
- $\bar{i} = \langle i_k : k \in \omega \rangle$ lists o.t. $(X \cap \alpha_*)$ without repetition in such a way that for each $k \in \omega$ the i_k th element of $X \cap \alpha_*$ is in Y_k ;
- $\bar{D} = \langle D_k : k \in \omega \rangle$ lists the dense open subsets of \mathbb{P}_{α_*} in X in such a way that each D_k is in the corresponding Y_k .

In the following definition, note that by the conditions on \bar{i} in Definition 3.2, α_{i_k} is in Y_k , for each $k \in \omega$.

{condtreedef}

3.3 Definition. Suppose that

$$S = \langle \bar{C}, F, \mathbf{q}, p, X, \langle Y_k : k \in \omega \rangle, \langle i_k : k \in \omega \rangle, \langle D_k : k \in \omega \rangle \rangle$$

is a suitable data sequence, and let

- \mathbf{q} be $\langle \mathbb{P}_\alpha, \dot{Q}_\beta, \dot{g}_\beta, \dot{h}_\beta : \alpha \leq \alpha_*, \beta < \alpha_* \rangle$,
- γ be $X \cap \omega_1$,
- $\bar{\alpha} = \langle \alpha_i : i < i_* \rangle$ list $X \cap \alpha_*$ in increasing order,
- \bar{C} be $\langle C_\alpha : \alpha \in \Omega \rangle$,
- for each $k \in \omega$, m_k be $|C_\gamma \cap Y_k|$ and u_k be $\{i_n : n < k\}$,
- α_{i_*} denote α_* and, for each $j \leq i_*$ and $k \in \omega$, $u_k(j)$ be $u_k \cap j$.

A *condition tree* for S is a pair $(T, \langle p_\rho : \rho \in T \rangle)$ such that

{itemone}

1. T is a finitely branching tree of finite sequences, with a unique node of length 1;

{itemtwo}

2. each $\rho \in T$ is sequence of the form $\langle s_k^\rho : k < |\rho| \rangle$ such that

{itemtwob}

- (a) each s_k^ρ is a function from u_k to 2^{m_k} ;
- (b) whenever $k + 1 < |\rho|$ and $i \in u_k$, $s_k^\rho(i) \sqsubseteq s_{k+1}^\rho(i)$;

{itemthree}

3. whenever $\rho \in T$ has length $k + 1$ and $s : u_k \rightarrow 2^{m_{k+1}}$ is such that, for all $i \in u_k$, $s_k^\rho(i) \sqsubseteq s(i)$, there is exactly one $\rho' \in T$ such that:

- $\rho \sqsubseteq \rho'$,
- $|\rho'| = k + 2$,
- $s \subseteq s_{k+1}^{\rho'}$;

{itemthreepfive}

4. whenever $\rho_1, \rho_2 \in T$ have length at least $k+2$ and $j < i_*$ is such that

$$s_{k+1}^{\rho_1} \upharpoonright u_k(j) = s_{k+1}^{\rho_2} \upharpoonright u_k(j),$$

then we have

$$s_{k+1}^{\rho_1} \upharpoonright u_{k+1}(j) = s_{k+1}^{\rho_2} \upharpoonright u_{k+1}(j);$$

{belp}

5. $p_{\langle \rangle} = p$;

{gooddec}

6. for all $\rho \in T \setminus \{\langle \rangle\}$, $p_\rho \in D_{|\rho|-1} \cap Y_{|\rho|-1}$;

{itemseven}

7. if $\rho_1 \trianglelefteq \rho_2 \in T$ then $p_{\rho_1} \geq_{\mathbb{P}_{\alpha_*}} p_{\rho_2}$;

8. if $\rho \in T$ and $|\rho| = k+1$ then for all $i \in u_k$,

$$p_\rho \upharpoonright (\alpha_i + 1) \Vdash_{\mathbb{P}_{\alpha_i+1}} \langle \dot{h}_{\alpha_i}(C_\gamma(n)) : n < m_k \rangle = \check{s}_k^\rho(i)$$

and $p_\rho \upharpoonright \alpha_i$ decides the value of $\delta_{p_\rho(\alpha_i)}$;

{itemeight}

9. if $\alpha \leq \alpha_*$ and $\rho_1, \rho_2 \in T$ have the same length and are such that

$$s_k^{\rho_1}(i) = s_k^{\rho_2}(i)$$

whenever $k < |\rho_1|$, $i \in u_k$ and $\alpha_i < \alpha$, then $p_{\rho_1} \upharpoonright \alpha = p_{\rho_2} \upharpoonright \alpha$.

3.4 Remark. Using parts (1) and (8) of Definition 3.3, the conjunction of parts (4) and (9) can be rephrased as the following statement : whenever ρ_1 and ρ_2 are distinct members of T of the same length $k+2$, letting i be the least member of u_{k+1} such that $s_{k+1}^{\rho_1}(i) \neq s_{k+1}^{\rho_2}(i)$, we have that $i \in u_k$ and α_i is the least α such that $p_{\rho_1} \upharpoonright (\alpha+1) \neq p_{\rho_2} \upharpoonright (\alpha+1)$.

The following lemma is an adaptation of the proof of Lemma 1.8 of [14].

{fusionlemma}

Lemma 3.5. *If S is a suitable data sequence, then there exists a condition tree relative to S .*

Proof. Let S be

$$\langle \bar{C}, F, \mathbf{q}, p, X, \langle Y_k : k \in \omega \rangle, \langle i_n : n < \omega \rangle, \langle D_k : k \in \omega \rangle \rangle,$$

and let

- γ be $X \cap \omega_1$,
- \mathbf{q} be $\langle \mathbb{P}_\alpha, \dot{Q}_\beta, \dot{g}_\beta, \dot{h}_\beta : \alpha \leq \alpha_*, \beta < \alpha_* \rangle$,
- $\bar{\alpha} = \langle \alpha_i : i < i_* \rangle$ list $X \cap \alpha_*$ in increasing order,
- \bar{C} be $\langle C_\alpha : \alpha \in \Omega \rangle$ and
- for each $k \in \omega$, m_k be $|C_\gamma \cap Y_k|$ and u_k be $\{i_n : n < k\}$.

We build T and $\langle p_\rho : \rho \in T \rangle$ by recursion on the length of $\rho \in T$. For each $k \in \omega$, let $T(k)$ denote the set of sequences in T of length k . For $k = 0$ we let $p_\langle \rangle = p$.

For each $k \in \omega$, let B_k be the (nonempty) set of conditions $q \leq p$ in $D_k \cap Y_k$ such that for each $i \in u_k$,

- $q \upharpoonright \alpha_i$ decides the value of $\delta_{q(\alpha_i)}$ (necessarily to be in $Y_k \cap \omega_1$);
- $q \upharpoonright (\alpha_i + 1)$ decides the value of $\langle \dot{h}_{\alpha_i}(C_\gamma(n)) : n < m_k \rangle$.

Given $q \in \mathbb{P}_{\alpha_*}$, $i < i_*$ and $k \in \omega$ such that q decides the value of the sequence $\langle \dot{h}_{\alpha_i}(C_\gamma(n)) : n < m_k \rangle$, let $t_{i,k}^q$ denote this decided value. Each $\rho \in T$ of length greater than zero will be determined by the corresponding p_ρ . That is, conditions (6) and (8) in Definition 3.3 imply that each p_ρ is in $B_{|\rho|-1}$ and each value $s_\ell^\rho(i)$ ($\ell < |\rho|$, $i \in u_\ell$) is the corresponding $t_{i,\ell}^{p_\rho}$.

To start, let q_1 be any member of B_0 . Let ρ_1 be the sequence of length 1 whose only member is the emptyset (noting that u_0 is the emptyset). Let $T(1) = \{\rho_1\}$ and let p_{ρ_1} be q_1 .

Suppose now that $k \in \omega \setminus \{0\}$ and that the members of $T(k)$ and the conditions $p_\rho \in B_{k-1}$ ($\rho \in T(k)$) have been chosen so as to satisfy the conditions in Definition 3.3. Let Z be the set of pairs (ρ, s) , where $\rho \in T(k)$ and $s : u_{k-1} \rightarrow 2^{m_k}$ are as in item (3) of Definition 3.3, that is, such that for all $i \in u_{k-1}$, $s_{k-1}^\rho(i) \leq s(i)$. Working in Y_k we pick for each pair $(\rho, s) \in Z$ a condition $p_{\rho,s} \leq p_\rho$ in B_k meeting conditions (8) and (9) of Definition 3.3 and such that the induced sequences $t_{i,k}^{p_{\rho,s}}$ ($i \in u_k$) will satisfy conditions (3) and (4) of Definition 3.3. Then the set of conditions corresponding to the members of $T(k+1)$ will be $\{p_{\rho,s} : (\rho, s) \in Z\}$, and this will induce the set $T(k+1)$.

We will do this in three stages. In the first stage we (eventually) choose conditions $p_{\rho,s}^{1, \max(u_k), |Z|} \leq p_\rho$ for each pair $(\rho, s) \in Z$, in such a way that the sequences $t_{i,k}^{p_{\rho,s}}$ are all determined (for each $(\rho, s) \in Z$ and $i \in u_k$, assuming, as will be the case, that the eventually chosen condition $p_{\rho,s}$ will be a strengthening of $p_{\rho,s}^{1, \max(u_k), |Z|}$). In the second stage we strengthen each condition $p_{\rho,s}^{1, \max(u_k), |Z|}$ to a condition $p_{\rho,s}^{2, |Z|}$ in D_k . In the third stage each condition $p_{\rho,s}^{2, |Z|}$ is strengthened to a condition $p_{\rho,s}^{3, \min(u_k), |Z|} = p_{\rho,s}$ deciding the value of $\delta_{p_{\rho,s}}(\alpha_i)$ for each $i \in u_k$. In each stage the choices are made simultaneously for all $(\rho, s) \in Z$ in order to ensure that conditions (4) and (9) of Definition 3.3 are preserved.

In the first stage we choose conditions of the form $p_{\rho,s}^{1, i, \ell} \leq p_\rho$ ($\ell \leq |Z|$, $i \in u_k$), recursively first in u_k (simultaneously for all $(\rho, s) \in Z$) and secondarily in $|Z|$ (although this is needed only the case $i = i_{k-1}$), in such a way that

{rhoscondone}

1. for all $(\rho, s) \in Z$,

(a) $p_{\rho,s}^{1, \min(u_k), 0} = p_\rho$;

{rhoscondonec}

(b) for all $i \in u_k$ and $\ell < |Z|$, $p_{\rho,s}^{1, i, \ell+1} \leq p_{\rho,s}^{1, i, \ell}$;

(c) for all $i \in u_k$, $\ell \leq |Z|$ and $\alpha \in \alpha_* \setminus (\alpha_i + 1)$, $p_{\rho,s}^{1, i, \ell}(\alpha) = p_\rho(\alpha)$;

(d) when $i < i'$ are successive members of u_k , $p_{\rho,s}^{1, i, |Z|} = p_{\rho,s}^{1, i', 0}$;

{rhoscondoneb}

(e) for each $i \in u_{k-1}$, $p_{\rho,s}^{1,i,1} = p_{\rho,s}^{1,i,|Z|}$ and

$$p_{\rho,s}^{1,i,1} \upharpoonright (\alpha_i + 1) \Vdash_{\mathbb{P}_{\alpha_i+1}} \langle \dot{h}_{\alpha_i}(C_\gamma(n)) : n < m_k \rangle = s(i);$$

{rhoscondoned}

(f) $p_{\rho,s}^{1,i_{k-1},|Z|} \upharpoonright (\alpha_{i_{k-1}} + 1)$ decides the value of

$$\langle \dot{h}_{\alpha_{i_{k-1}}}(C_\gamma(n)) : n < m_k \rangle.$$

{rhoscondtwo}

2. for all $(\rho_1, s_1), (\rho_2, s_2) \in Z$, all $i \in u_k$ all $\ell \leq |Z|$ and all $\alpha \leq \alpha_*$, if

$$s_1 \upharpoonright \{j \in u_{k-1} : \alpha_j < \alpha\} = s_2 \upharpoonright \{j \in u_{k-1} : \alpha_j < \alpha\}$$

$$\text{then } p_{\rho_1, s_1}^{1,i,\ell} \upharpoonright \alpha = p_{\rho_2, s_2}^{1,i,\ell} \upharpoonright \alpha.$$

Note that item (2) above corresponds to condition (9) of Definition 3.3, which holds for the conditions p_ρ for $\rho \in T(k)$ by the recursive hypothesis.

We choose the conditions $p_{\rho,s}^{1,i,\ell}$ ($(\rho, s) \in Z$, $1 \leq \ell \leq |Z|$) as follows. For each $(\rho, s) \in Z$, let $p_{\rho,s}^{1,i,0}$ be p_ρ if $i = \min(u_k)$. Otherwise let $p_{\rho,s}^{1,i,0}$ be $p_{\rho,s}^{1,i',|Z|}$, where i' is the largest member of u_k below i . Enumerate $|Z|$ as $\langle (\rho_\ell, s_\ell) : \ell < |Z| \rangle$.

First consider the case $i \in u_{k-1}$. Define an equivalence relation E_i on Z consisting of the pairs $((\rho, s), (\rho', s'))$ such that

$$s \upharpoonright \{j \in u_{k-1} : \alpha_j \leq \alpha_i\} = s' \upharpoonright \{j \in u_{k-1} : \alpha_j \leq \alpha_i\}.$$

By the recursive hypothesis applied to item (2), for any two E_i -equivalent pairs (ρ, s) and (ρ', s') in Z ,

$$p_{\rho,s}^{1,i,0} \upharpoonright (\alpha_i + 1) = p_{\rho',s'}^{1,i,0} \upharpoonright (\alpha_i + 1).$$

Each condition $p_{\rho,s}^{1,i,0}$ is below p_ρ , which forces that $\delta_{p_\rho(\alpha_i)} < Y_{k-1} \cap \omega_1$. Furthermore, $Y_{k-1} \cap \omega_1$ is below the least element of C_γ for which the value of \dot{h}_{α_i} is not decided by p_ρ . By item (1c) (applied to the previous values of i , if there are any), $p_{\rho,s}^{1,i,0}(\alpha_i) = p_\rho(\alpha_i)$ for all $(\rho, s) \in Z$. We can then strengthen each condition $p_{\rho,s}^{1,i,0}$ ($(\rho, s) \in Z$) to a condition $p_{\rho,s}^{1,i,1}$ while changing only coordinate α_i (in the same way for E_i -equivalent pairs) to make the displayed formula from item (1e) hold. As in item (1e), for all $\ell \in (1, |Z|]$, we let $p_{\rho,s}^{1,i,\ell} = p_{\rho,s}^{1,i,1}$.

When $i \in u_k \setminus u_{k-1}$ (i.e., $i = i_{k-1}$), our task is easier in that there are no previously specified values of $\langle \dot{h}_{\alpha_i}(C_\gamma(n)) : n < m_k \rangle$ to obtain. However, it is harder in that we may need to change our conditions in coordinates below α_i while preserving item (2) at lower values of α (conditions disagreeing on the interval $[0, \alpha_i]$ may still agree on some such interval $[0, \alpha]$). Our method for doing this will be used again in stages two and three.

For each $\ell < |Z|$, given the conditions $p_{\rho,s}^{1,i,\ell}$ we strengthen condition $p_{\rho_\ell, s_\ell}^{1,i,\ell}$ (perhaps trivially, and only in coordinates less than or equal to α_i) to decide the value of $\langle \dot{h}_{\alpha_i}(C_\gamma(n)) : n < m_k \rangle$, thus forming condition $p_{\rho_\ell, s_\ell}^{1,i,\ell+1}$ (still in Y_k).

For each other condition $p_{\rho,s}^{1,i,\ell}$ (i.e., with $(\rho, s) \in Z \setminus \{(\rho_\ell, s_\ell)\}$), letting α be maximal with

$$s \upharpoonright \{j \in u_{k-1} : \alpha_j < \alpha\} = s_\ell \upharpoonright \{j \in u_{k-1} : \alpha_j < \alpha\},$$

(by item (2) this implies that $p_{\rho,s}^{1,i,\ell} \upharpoonright \alpha = p_{\rho_\ell, s_\ell}^{1,i,\ell} \upharpoonright \alpha$), we let $p_{\rho,s}^{1,i,\ell+1}$ be the concatenation of $p_{\rho_\ell, s_\ell}^{1,i,\ell+1} \upharpoonright \alpha$ with $p_{\rho,s}^{1,i,\ell} \upharpoonright [\alpha, \alpha_*)$.

The choice of the conditions $p_{\rho,s}^{1,\max(u_k),|Z|}$ induces the corresponding members of $T(k+1)$: letting, for each $(\rho, s) \in Z$, $t_{\rho,s} : u_k \rightarrow 2^{m_k}$ be such that

$$t_{\rho,s}(i) = t_{i,k}^{p_{\rho,s}^{1,\max(u_k),|Z|}},$$

we let $T(k+1)$ be

$$\{\rho \widehat{\ } \langle t_{\rho,s} \rangle : (\rho, s) \in Z\}.$$

The members of $T(k+1)$ have been chosen to satisfy conditions (3) and (4) of Definition 3.3.

In stages 2 and 3 we will strengthen each $p_{\rho,s}^{1,\max(u_k),|Z|}$ to a condition $p_{\rho,s}$ by modifications of the argument above for item (1f). We will then let $p_{\rho,s}$ be $p_{\rho \widehat{\ } \langle t_{\rho,s} \rangle}$ for each $(\rho, s) \in Z$. Condition (1e) above implies that any strengthening of any $p_{\rho,s}^{1,\max(u_k),|Z|}$ will satisfy condition (8) of Definition 3.3, aside from the requirement that the values of $\delta_{p_{\rho,s}(\alpha_i)}$ ($i \in u_k$) must be decided, which we will take care of in stage 3.

The recursive hypotheses and item (2) above imply that the conditions $p_{\rho,s}^{1,\max(u_k),|Z|}$ do not violate condition (9) of Definition 3.3. In particular, we have that for all distinct pairs $(\rho, s), (\rho', s')$ from Z , the greatest $\alpha \leq \alpha_*$ for which $p_{\rho,s}^{1,\max(u_k),|Z|} \upharpoonright \alpha = p_{\rho',s'}^{1,\max(u_k),|Z|} \upharpoonright \alpha$ is also the least value α_i ($i \in u_{k-1}$) for which $s(i) \neq s'(i)$. We will maintain this equivalence throughout the rest of the construction for the various conditions associated to the pairs (ρ, s) from Z .

We now (in stage 2) strengthen each $p_{\rho,s}^{1,\max(u_k),|Z|}$ to a condition $p_{\rho,s}^{2,|Z|}$ in D_k , while respecting condition (9) of Definition 3.3. We reuse our enumeration of $|Z|$ as $\langle (p_\ell, s_\ell) : \ell < |Z| \rangle$. We choose conditions $p_{\rho,s}^{2,\ell}$ for $(\rho, s) \in Z$ and $\ell \leq |Z|$, recursively in ℓ so that:

3. for all $(\rho, s) \in Z$,

$$(a) \ p_{\rho,s}^{2,0} = p_{\rho,s}^{1,\max(u_k),|Z|},$$

$$(b) \ \text{for all } \ell < |Z|, \ p_{\rho,s}^{2,\ell+1} \leq p_{\rho,s}^{2,\ell} \text{ and } p_{\rho_\ell, s_\ell}^{2,\ell+1} \in D_k;$$

{coherencec}

4. for all $\ell \leq |Z|$, and all distinct $(\rho, s), (\rho', s') \in Z$, the least α such that

$$p_{\rho,s}^{2,\ell}(\alpha) \neq p_{\rho',s'}^{2,\ell}(\alpha)$$

is the same as the least α such that

$$p_{\rho,s}^{2,0}(\alpha) \neq p_{\rho',s'}^{2,0}(\alpha).$$

Item (4) ensures the preservation of condition (9) of Definition 3.3, i.e., that, for all $(\rho, s), (\rho', s') \in Z$ and all $\alpha \leq \alpha_*$, if

$$s \upharpoonright \{j \in u_k : \alpha_j < \alpha\} = s' \upharpoonright \{j \in u_k : \alpha_j < \alpha\}$$

then $p_{\rho, s}^{2, \ell} \upharpoonright \alpha = p_{\rho', s'}^{2, \ell} \upharpoonright \alpha$.

In substage ℓ of stage 2, strengthen $p_{\rho_\ell, s_\ell}^{2, \ell}$ (possibly trivially) to a condition $p_{\rho_\ell, s_\ell}^{2, \ell+1}$ in D_k . For each of the other conditions $p_{\rho, s}^{2, \ell}$, letting α be minimal such that

$$p_{\rho_\ell, s_\ell}^{2, \ell}(\alpha) \neq p_{\rho, s}^{2, \ell}(\alpha)$$

let $p_{\rho, s}^{2, \ell+1}$ be the concatenation of $p_{\rho_\ell, s_\ell}^{2, \ell+1} \upharpoonright \alpha$ with $p_{\rho, s}^{2, \ell} \upharpoonright [\alpha, \alpha_*)$.

Finally, in stage 3, working recursively through the members of u_k from top to bottom, and still working in Y_k , we strengthen each $p_{\rho, s}^{2, |Z|}$ to a condition $p_{\rho, s}$ which decides each value $\delta_{p_{\rho, s}(\alpha_i)}$ ($i \in u_k$), while preserving item (2) from earlier in this proof and thus condition (9) from Definition 3.3. For each $i \in u_k$ we can accomplish this in $|Z|$ many steps via an argument similar to the ones used in stages 1 and 2. We now choose conditions $p_{\rho, s}^{3, i, \ell}$ for $(\rho, s) \in Z$, $i \in u_k$, $\ell \leq |Z|$ such that

5. for all $(\rho, s) \in Z$,

(a) $p_{\rho, s}^{3, \max(u_k), 0} = p_{\rho, s}^{2, |Z|}$;

(b) for all $i \in u_k$ and $\ell < |Z|$, $p_{\rho, s}^{3, i, \ell+1} \leq p_{\rho, s}^{3, i, \ell}$;

(c) for all $i \in u_k$, $\ell \leq |Z|$ and $\alpha \in \alpha_* \setminus \alpha_i$, $p_{\rho, s}^{3, i, \ell+1}(\alpha) = p_{\rho, s}^{3, i, \ell}(\alpha)$;

(d) when $i' < i$ are successive members of u_k , $p_{\rho, s}^{3, i, |Z|} = p_{\rho, s}^{3, i', 0}$;

(e) for all $i \in u_k$ and $\ell < |Z|$, $p_{\rho, s}^{3, i, \ell+1} \upharpoonright \alpha_i$ decides the value of $\delta_{p_{\rho, s}^{3, i, \ell+1}(\alpha_i)}$;

6. for all $i \in u_k$ and $\ell \leq |Z|$, and all distinct $(\rho, s), (\rho', s') \in Z$, the least α such that

$$p_{\rho, s}^{3, i, \ell}(\alpha) \neq p_{\rho', s'}^{3, i, \ell}(\alpha)$$

is the same as the least α such that

$$p_{\rho, s}^{2, 0}(\alpha) \neq p_{\rho', s'}^{2, 0}(\alpha).$$

In the substage of stage 3 corresponding to $i \in u_k$ and $\ell < |Z|$, strengthen $p_{\rho_\ell, s_\ell}^{3, i, \ell}$ (possibly trivially, and only in coordinates below α_i) to a condition $p_{\rho_\ell, s_\ell}^{3, i, \ell+1}$ deciding the value of $\delta_{p_{\rho_\ell, s_\ell}^{3, i, \ell+1}(\alpha_i)}$. For each of the other conditions $p_{\rho, s}^{3, i, \ell}$, letting α be minimal such that

$$p_{\rho_\ell, s_\ell}^{3, i, \ell}(\alpha) \neq p_{\rho, s}^{3, i, \ell}(\alpha)$$

let $p_{\rho, s}^{3, i, \ell+1}$ be the concatenation of $p_{\rho_\ell, s_\ell}^{3, i, \ell+1} \upharpoonright \alpha$ with $p_{\rho, s}^{3, i, \ell} \upharpoonright [\alpha, \alpha_*)$.

Finally, for each $(\rho, s) \in Z$ let $p_{\rho \frown \langle t_{\rho, s} \rangle} = p_{\rho, s}$ be $p_{\rho, s}^{3, \min(u_k), |Z|}$. \square

We will need to consider restrictions of our condition trees to initial segments of our iterations. Given a tree T in \mathcal{T} , we let $[T]$ denote the set of infinite branches through T .

{projdefs}

3.6 Definition. Suppose that $(T, \langle p_\rho : \rho \in T \rangle)$ is a condition tree relative to some suitable data sequence

$$S = \langle \bar{C}, F, \mathbf{q}, p, X, \langle Y_k : k < \omega \rangle, \langle i_n : n < \omega \rangle, \bar{D} \rangle,$$

$\gamma = X \cap \omega_1$, α_* is the length of \mathbf{q} , and that $\langle \alpha_i : i < i_* \rangle$ enumerates $X \cap \alpha_*$ in increasing order. For each $k \in \omega$, let $u_k = \{i_n : n < k\}$. Let α_{i_*} denote α_* .

- For each $j \leq i_*$, let
 - for each $k \in \omega$, $u_k(j)$ be $u_k \cap j$;
 - for each $\rho \in T$, $\rho^{[j]}$ be

$$\langle s_\ell^\rho \upharpoonright u_\ell(j) : \ell < |\rho| \rangle,$$

- let $s_\ell^{\rho^{[j]}}$ (for $\ell < |\rho^{[j]}|$) denote $s_\ell^\rho \upharpoonright u_\ell(j)$, and $p_\rho^{[j]}$ be $p_\rho \upharpoonright \alpha_j$;
- $T^{[j]}$ be the tree of sequences $\{\rho^{[j]} : \rho \in T\}$.

- For all $i \leq j < i_*$ and each $\rho \in T^{[j]}$, let $\rho^{[i]}$ be $\langle \rho(\ell) \upharpoonright u_\ell(i) : \ell < |\rho| \rangle$.
- For all $i \leq j \leq i_*$, let
 - $\text{proj}_{j,i} : T^{[j]} \rightarrow T^{[i]}$ be the function defined by setting

$$\text{proj}_{j,i}(\rho) = \rho^{[i]};$$

- for each $x \in [T^{[i]}]$, $T^{[j]}(x)$ be $\{\rho \in T^{[j]} : \rho^{[i]} \leq x\}$;
- for each $x \in [T^{[j]}]$, $x^{[i]}$ be the union of $\{(x \upharpoonright k)^{[i]} : k \in \omega\}$.

We record some observations on these definitions.

{computationthrm}

Theorem 3.7. Suppose that $(T, \langle p_\rho : \rho \in T \rangle)$ is a condition tree relative to some suitable data sequence

$$S = \langle \bar{C}, F, \mathbf{q}, p, X, \langle Y_k : k < \omega \rangle, \langle i_n : n < \omega \rangle, \bar{D} \rangle,$$

$\gamma = X \cap \omega_1$, α_* is the length of \mathbf{q} , and that $\langle \alpha_i : i < i_* \rangle$ enumerates $X \cap \alpha_*$ in increasing order. For each $\ell \in \omega$, let $u_\ell = \{i_n : n < \ell\}$. Let α_{i_*} denote α_* .

{remtwo}

1. For all $\rho \in T$ and $\ell \in \omega$, $\rho^{[i_*]} = \rho$ and $u_\ell(i_*) = u_\ell$.

{remthree}

2. If $j \leq i_*$ and $\rho \in T^{[j]}$ has length $k+1$, then ρ has exactly $2^{|u_k(j)|(m_{k+1}-m_k)}$ many immediate successors in $T^{[j]}$.

3. For all $i \leq j \leq i_*$ and $x \in [T^{[i]}]$, if $\rho \in T^{[j]}(x)$ has length $k+1$, then ρ has exactly $2^{|u_k(j) \setminus u_k(i)|(m_{k+1}-m_k)}$ many successors in $T^{[j]}(x)$.

{remfour}

4. For all $i \leq j \leq i_*$, $k \in \omega \setminus \{0\}$, $\rho \in T^{[i]}$ of length $k + 1$ and $\nu \in T^{[j]}$ of length k with $\nu^{[i]} = \rho \upharpoonright k$, the set

$$\{\tau \in T^{[j]} : \tau \upharpoonright k = \nu, \tau^{[i]} = \rho\}$$

{remfive}

has size $2^{|u_{k-1}(j) \setminus u_{k-1}(i)|(m_k - m_{k-1})}$.

5. For all $c \leq i \leq j \leq i_*$, $x \in [T^{[c]}]$, $k \in \omega \setminus \{0\}$, $\rho \in T^{[i]}(x)$ of length $k + 1$ and $\nu \in T^{[j]}(x)$ of length k with $\nu^{[i]} = \rho \upharpoonright k$, the set

$$\{\tau \in T^{[j]}(x) : \tau \upharpoonright k = \nu, \tau^{[i]} = \rho\}$$

has size $2^{|u_{k-1}(j) \setminus u_{k-1}(i)|(m_k - m_{k-1})}$.

Proof. The first conclusion follows immediately from the definitions. For part (2), note that by parts (2b), (3) and (4) of Definition 3.3, the immediate successors of a $\rho \in T^{[j]}$ of length $k + 1$ correspond to the functions from $u_k(j)$ to $2^{m_{k+1}}$ which pointwise extend s_k^ρ , of which there are $2^{|u_k(j)|(m_{k+1} - m_k)}$ many. For part (2), the same parts of Definition 3.3 imply that the immediate successors of a $\rho \in T^{[j]}(x)$ of length $k + 1$ correspond to the functions from $u_k(j) \setminus u_k(i)$ to $2^{m_{k+1}}$ which pointwise extend $s_k^\rho \upharpoonright (u_k(j) \setminus u_k(i))$, of which there are $2^{|u_k(j) \setminus u_k(i)|(m_{k+1} - m_k)}$ many.

For part (4), the immediate successors of ν correspond to the functions $s: u_{k-1}(j) \rightarrow 2^{m_k}$ which pointwise extend s_{k-1}^ν , and the successors τ of ν with $\tau^{[i]} = \rho$ correspond to those s which agree with s_k^ρ on $u_{k-1}(i)$. The number of such functions is $2^{|u_{k-1}(j) \setminus u_{k-1}(i)|(m_k - m_{k-1})}$. Part (5) is similar. \square

3.8 Remark. Using the objects introduced in the statement of Theorem 3.7, fix for each pair (i, j) with $i < j \leq i_*$ a tree $T_{i,j}$ of finite sequences with unique nodes of lengths 0 and 1 and each node of each length $k > 0$ having exactly $2^{|u_{k-1}(j) \setminus u_{k-1}(i)|(m_k - m_{k-1})}$ many immediate successors. The computations in Theorem 3.7 show that each tree $T^{[j]}$ is isomorphic to the tree of pairs of sequences (ρ, ν) of the same length with $\rho \in T^{[i]}$ and $\nu \in T_{i,j}$. Similarly, each tree $T^{[j]}(x)$ for some $x \in [T^{[c]}]$ for some $c \leq i$ is isomorphic to the tree of pairs of sequences (ρ, ν) of the same length with $\rho \in T^{[i]}(x)$ and $\nu \in T_{i,j}$. Under these identifications, the function $\text{proj}_{j,i}$ from Definition 3.6 corresponds to the usual first-coordinate projection function. These facts let us apply Fubini's theorem and the Kuratowski-Ulam theorem to pairs $[T^{[i]}], [T^{[j]}]$ (with the latter viewed as $[T^{[i]}] \times [T_{i,j}]$) and pairs $[T^{[i]}(x)], [T^{[j]}(x)]$ (with the latter viewed as $[T^{[i]}(x)] \times [T_{i,j}]$), which we do in the proof of Lemma 5.4.

{tclosedrem}

The following observation will be used at the end of the proof of Lemma 5.4.

3.9 Remark. Suppose that $(T, \langle p_\rho : \rho \in T \rangle)$ is a condition tree for a suitable data sequence $S = \langle \bar{C}, F, \mathbf{q}, p, X, Y, \bar{1}, \bar{D} \rangle$, and that i_* is the ordertype of $X \cap \alpha_*$, where α_* is the ordertype of the iteration \mathbf{q} . Let $j \leq i_*$ and let K be a closed subset of $[T^{[j]}]$. Then for all $x \in [T^{[j]}]$, $x \in K$ if and only if, for cofinally many $c < j$, $x^{[c]} \in \text{proj}_{j,c}[K]$.

4 Preserving pathology

{pathsec}

In this section we add a condition on our dense partial function F , and recall some important properties of the null and meager ideals. We say that an ideal I on a Polish space is *Borel* if it is generated by Borel sets (i.e., every member of the ideal is contained in a Borel member of the ideal). The meager and null ideals are Borel in this sense. Given an ideal I on a set X , we let I^+ denote $\mathcal{P}(X) \setminus I$, and we say that a subset of X is *I -Borel-large* if it intersects every Borel set in I^+ .

4.1 Definition. Given a topological space X , an ideal I on X , and a set S with at least two elements, we say that a partial function $F: X \rightarrow S$ is *I -pathological* if for every $s \in S$, $F^{-1}[\{s\}]$ is I -Borel-large.

When I is the ideal of countable sets, we say that F is *totally pathological*. We say that F is *Lebesgue-pathological* when I is the ideal of Lebesgue null sets, and *category-pathological* when I is the ideal of meager sets. Note that total pathology implies both Lebesgue and category pathology.

4.2 Remark. A totally pathological function is dense in the sense of Definition 2.1, and the existence of a totally pathological function implies the existence of sets of reals without the perfect set property.

A standard construction, using a wellordering of the continuum and the fact that uncountable Borel sets have cardinality continuum, shows that ZFC implies the existence of (total) totally pathological functions on 2^ω . Moreover, if a is a subset of ω , then the same construction, using a $\Sigma_2^1(a)$ wellordering of $(2^\omega)^{\mathbf{L}[a]}$, shows that there is in $\mathbf{L}[a]$ a total totally pathological function $F: 2^\omega \rightarrow 2$ which is Σ_2^1 in a .

In our proof in Section 5 we iterate forcings of the form $Q_{\bar{C},F,g}$ using a fixed function F which is totally pathological and $\Sigma_2^1(a)$ in the ground model $\mathbf{L}[a]$. Since our iterations will add reals, the function F will not remain totally pathological in the corresponding forcing extensions. For the proof of our main theorem, we need to know that F remains I -pathological throughout the iteration, where I is either the null ideal or the meager ideal.

{prespropdef}

4.3 Definition. A Borel ideal I has the *preservation property* if every I -Borel-large set remains so after any countable support iteration of partial orders of the form $Q_{\bar{C},F,g}$.

{pathpresrem}

4.4 Remark. The ideals that we consider in this paper are ideals on the set of infinite branches through finitely branching trees of height ω . Theorem 5.2 of [8] implies that the versions of the meager ideal on these spaces have the preservation property (for a much wider class of partial orders than the ones considered here). Theorem 6.3 and Lemma 6.4 of [8] show the same thing for the ideals of Lebesgue null sets.

The proof of Theorem 5.1 will also use the fact that the meager and null ideals are regular, in the sense that every Borel set not in either ideal contains a closed set which is not in the same ideal.

5 Measure and Category

{itsec}

In this section we prove our main theorem.

{mainthrm}

Theorem 5.1. *Suppose that a is a subset of ω such that $\mathbf{V} = \mathbf{L}[a]$, and let \bar{C}_a be the canonical ladder system relative to a . Let F be a totally pathological dense function whose graph is Σ_2^1 in $\mathbf{L}[a]$. Suppose that*

$$\langle \mathbb{P}_\alpha, \dot{Q}_\beta, \dot{g}_\beta, \dot{h}_\beta : \alpha \leq \omega_2^{\mathbf{V}}, \beta < \omega_2^{\mathbf{V}} \rangle$$

is a fully bookkeeping element of $\mathbf{Q}_{\bar{C}_a, F}$. Then $\mathbb{P}_{\omega_2^{\mathbf{V}}}$ forces that every subset of ω^ω which is either universally measurable or universally categorical is Δ_2^1 .

The proof of Theorem 5.1 runs from here through the end of this section. We will concentrate on the universally measurable case, and note the changes that need to be made for universally categorical sets.

By Remark 1.4, it suffices to prove that each tail of any iteration as in the statement of Theorem 5.1 is \mathcal{A} -representing, where \mathcal{A} is either the collection of universally measurable subsets of ω^ω or the collection of universally categorical sets. Since the ideals of Lebesgue-null sets and meager sets have the preservation property (as in Remark 4.4), in each intermediate model of such a forcing iteration, F is Lebesgue-pathological and category-pathological (but no longer totally pathological once the iteration has added new subsets of ω). It remains then to prove the following.

{representinglem}

Lemma 5.2. *Suppose that*

- \bar{C} is a ladder system on ω_1 ,
- $F: 2^\omega \rightarrow 2$ is a dense function which is Lebesgue-pathological and category-pathological,
- $\mathbf{q} = \langle \mathbb{P}_\alpha, \dot{Q}_\beta, \dot{g}_\beta, \dot{h}_\beta : \alpha \leq \alpha_*, \beta < \alpha_* \rangle$ is in $\mathbf{Q}_{\bar{C}, F}$ and
- $\bar{\mathbb{P}}$ is forcing iteration $\langle \mathbb{P}_\alpha, \dot{Q}_\beta : \alpha \leq \alpha_*, \beta < \alpha_* \rangle$.

Then \mathbb{P}_{α_} is \mathcal{A} -representing, where \mathcal{A} is either the collection of universally measurable subsets of ω^ω or the collection of universally categorical sets.*

Fix the objects introduced in the statement of Lemma 5.2 from here through the end of the proof of Lemma 5.4, which will complete the proof of Lemma 5.2 and thus Theorem 5.1. We have to show that if

- p is a condition in \mathbb{P}_{α_*} ,
- $A \subseteq \omega^\omega$ is either universally measurable or universally categorical,
- τ is a \mathbb{P}_{α_*} -name for an element of ω^ω

then there exist a condition $q \leq p$ and a Borel set $B \subseteq \omega^\omega$ such that

- B is either contained in or disjoint from A and
- $q \Vdash \tau \in \check{B}$.

In fact the set we find will be a continuous image of a Borel set; as every analytic set is a union of \aleph_1 many Borel sets (in an absolute way, see page 201 of [7]), this suffices. To find our desired condition and analytic set, fix such p , A and τ , and fix in addition X , \bar{Y} , $\bar{\iota}$ and \bar{D} (with $\tau \in X$) such that $S = \langle \bar{C}, F, \mathbf{q}, p, X, \bar{Y}, \bar{\iota}, \bar{D} \rangle$ is a suitable data sequence. By Lemma 3.5, we may fix in addition a condition tree $(T, \langle p_\rho : \rho \in T \rangle)$ relative to S . There is then a continuous function $f: [T] \rightarrow \omega^\omega$ such that each value $f(x)$ is the realization of τ by $\{p_{x \upharpoonright n} : n \in \omega\}$, since any such set meets all the dense open subsets of \mathbb{P}_{α_*} in X . Since A is universally measurable (or universally categorical), there exist then Borel sets B and N contained in $[T]$ such that N is Lebesgue null (or meager) and $f^{-1}[A] \triangle B \subseteq N$. We will find a condition $p' \leq p$ forcing the existence (in the forcing extension) of an $x \in [T] \setminus N$ (reinterpreted) such that $\{p_{x \upharpoonright n} : n \in \omega\}$ is a subset of the generic filter. Some condition $q \leq p'$ will then decide whether the realization of τ is in the reinterpretation of $f[B]$ or not. In the former case $f[B \setminus N]$ is the desired analytic set; in the latter $f[[T] \setminus (B \cup N)]$ is. So our remaining task is just to find p' as above.

We will in fact start with a closed non-null (or nonmeager) set $K \subseteq [T]$ disjoint from N and find a condition $p' \leq p$ forcing the existence of an $x \in K$ (reinterpreted) such that $\{p_{x \upharpoonright n} : n \in \omega\}$ is a subset of the generic filter. As in the previous paragraph, this will complete the proofs of Lemma 5.2 and Theorem 5.1. An instance of such a condition will be a Lebesgue- (i_*, \check{K}) -solution (respectively, a category- (i_*, \check{K}) -solution) relative to S and $(T, \langle p_\rho : \rho \in T \rangle)$, as in Definition 5.3 below.

Given $j \leq i_*$, we let $\dot{\sigma}_j$ be a \mathbb{P}_j -name for the set of $\rho \in T^{[j]}$ for which $p_\rho \in G_j$, where G_j denotes the restriction of the generic filter G to \mathbb{P}_{α_j} . In the desired case where this set is infinite, we identify it with the corresponding element of $[T^{[j]}]$.

{soldef}

5.3 Definition. Suppose that $(T, \langle p_\rho : \rho \in T \rangle)$ is a condition tree relative to some suitable data sequence

$$S = \langle \bar{C}, F, \mathbf{q}, p, X, \langle Y_k : k \in \omega \rangle, \langle i_n : n < \omega \rangle, \bar{D} \rangle.$$

Let

- \mathbf{q} be $\langle \mathbb{P}_\alpha, \dot{Q}_\beta, \dot{g}_\beta, \dot{h}_\beta : \alpha \leq \alpha_*, \beta < \alpha_* \rangle$;
- $\langle \alpha_i : i < i_* \rangle$ enumerate $X \cap \alpha_*$ in increasing order;
- α_{i_*} denote α_* ;
- j be an element of $i_* + 1$;
- \dot{K} be a \mathbb{P}_j -name for a subset of $[T]$;

- q be a condition in \mathbb{P}_{α_j} ;

We say that q is a Lebesgue- (j, \dot{K}) -solution if q forces that

- $\dot{\sigma}_j \in [T^{[j]}]$;
- if $j < i_*$ then $\dot{K} \cap [T(\dot{\sigma}_j)]$ is non-null in $[T(\dot{\sigma}_j)]$;
- if $j = i_*$ then $\dot{\sigma}_j \in \dot{K}$.

We say that q is a category- (j, \dot{K}) -solution if q forces that

- $\dot{\sigma}_j \in [T^{[j]}]$;
- if $j < i_*$ then $\dot{K} \cap [T(\dot{\sigma}_j)]$ is nonmeager in $[T(\dot{\sigma}_j)]$;
- if $j = i_*$ then $\dot{\sigma}_j \in \dot{K}$.

The statement of the following lemma uses the objects introduced in this section. The case where $i = 0$, $j = i_*$ and $\dot{K} = \check{K}$ for some non-null closed K disjoint from N as above proves Lemma 5.2 and thereby completes the proof of Theorem 5.1.

{finallemma}

Lemma 5.4. *Suppose that $i < j \leq i_*$, \dot{K} is a \mathbb{P}_i -name for a closed subset of $[T]$ and q is a Lebesgue- (i, \dot{K}) -solution (respectively, a category- (i, \dot{K}) -solution). Then there is a Lebesgue- (j, \dot{K}) -solution (category- (j, \dot{K}) -solution) $q' \in \mathbb{P}_{\alpha_j}$ such that $q' \upharpoonright \alpha_i = q$.*

Proof. We prove this by induction on j for all i and \dot{K} simultaneously. We first note that the case $j = j' + 1$ reduces to the case where $j = i + 1$ by the induction hypothesis.

For the case where $j = i + 1$, suppose that $G_i \subseteq \mathbb{P}_{\alpha_i}$ is a generic filter with $q \in G_i$. Then $G_i \cap X$ is X -generic. Let e be $\dot{g}_{\alpha_i, G_i}(X \cap \omega_1)$, let $y = \dot{\sigma}_{i, G_i}$ and let \hat{y} be the function from i to 2^ω induced by y , so that, for all $c < i$, $\hat{y}(c) = \bigcup \{s_k^\rho(c) : \rho \in y, k < |\rho|, c \in u_k\}$. Applying Fubini's theorem (or the Kuratowski-Ulam theorem) and the regularity of the null ideal (or meager ideal), let K' be a closed non-null (nonmeager) subset of $[T^{[j]}(y)]$ with the property that for each $z \in K'$ the set $\{x \in K_{G_i} : \text{proj}_{i_*, j}(x) = z\}$ is non-null (nonmeager).

Let $k \in \omega$ be minimal with $i \in u_k$. Then $k > 0$. Since i is not in u_{k-1} , the tree $T^{[j]}(y)$ has a unique node ρ_k of length k , where for each $\ell < k$ and each $c \in u_\ell(j)$, c is in $u_\ell(i)$ and $s_\ell^{\rho_k}(c) = \hat{y}(c) \upharpoonright m_\ell$. Moreover, for all ρ, ρ' of length $k + 1$ in $T(y)$ (formally defined as $T^{[i_*]}(y)$), $s_k^\rho \upharpoonright u_{k-1}(j) = s_k^{\rho'} \upharpoonright u_{k-1}(j)$ (as just established), so, by condition (4) of Definition 3.3, $s_k^{\rho'} \upharpoonright u_k(j) = s_k^{\rho'} \upharpoonright u_k(j)$. It follows that $T^{[j]}(y)$ has a unique node ρ_{k+1} of length $k + 1$. The condition $\rho_{k+1} \upharpoonright \alpha_j$ forces

$$\langle \dot{h}_{\alpha_i}(C_{X \cap \omega_1}(n)) : n < m_k \rangle$$

to be $s_k^{\rho_{k+1}}(i)$, which is in 2^{m_k} . For each $\ell \geq k + 1$, each node ρ of $T^{[j]}(y)$ of length ℓ has $2^{m_\ell - m_{\ell-1}}$ many immediate successors, corresponding to the possible extensions of $s_{\ell-1}^\rho(i)$ of length m_ℓ .

Let T_* be the tree of elements of $2^{<\omega}$ compatible with the sequence $s_k^{\rho_{k+1}}(i)$. Thus $[T_*]$ is a clopen subset of 2^ω , and equal to

$$\{\bigcup\{s_\ell^{z \upharpoonright (\ell+1)}(i) : \ell \in \omega \setminus k\} : z \in [T^{[j]}(y)]\}.$$

By Remark 4.4, F is both Lebesgue-pathological and category-pathological in $V[G_i]$. Since the null (meager) ideal on $[T_*]$ is isomorphic to that on $[T^{[j]}(y)]$ there is a $z \in K'$ such that

$$F \left(\bigcup_{\ell \in [k, \omega)} s_\ell^{z \upharpoonright (\ell+1)}(i) \right) = e.$$

We can then let q' be (q, \dot{r}) , where \dot{r} is a \mathbb{P}_{α_i} name for $\bigcup\{s_\ell^{z \upharpoonright (\ell+1)}(i) : \ell \in \omega \setminus k\}$, which is an $X[G_i \cap X]$ -generic condition in \dot{Q}_{α_i, G_i} .

For the case where j is a limit ordinal, applying Fubini's theorem (or the Kuratowski-Ulam theorem) again, let \dot{K}_0 be a \mathbb{P}_i -name for a closed subset of (the realization of) \dot{K} such that q is a Lebesgue- (i, \dot{K}_0) -solution (category- (i, \dot{K}_0) -solution) and, q forces in addition that for all $y \in \text{proj}_{i_*, j}[\dot{K}_0]$ the set $\dot{K} \cap [T(y)]$ is non-null (nonmeager) (of course, this step is not necessary if $j = i_*$). This case now follows from the induction hypothesis applied successively to \dot{K}_0 and the members of any cofinal ω -sequence in the interval (i, j) , along with Remark 3.9. \square

6 Borel reinterpretations

Finally, we show that the Borel reinterpretations of universally measurable sets under the forcing extensions considered here are again universally measurable.

{umeasreint}

Theorem 6.1. *Let \bar{C} be a ladder system on ω_1 and let $F: 2^\omega \rightarrow 2$ be a null-pathological dense partial function. Let*

$$\langle \mathbb{P}_\alpha, \dot{Q}_\beta, \dot{g}_\beta, \dot{h}_\beta : \alpha \leq \alpha_*, \beta < \alpha_* \rangle,$$

be an element of $\mathbf{Q}_{\bar{C}, F}$. If $A \subseteq \omega^\omega$ is universally measurable, then the Borel reinterpretation of A is universally measurable in any forcing extension by \mathbb{P}_{α_} .*

Proof. It follows from Lemma 5.2 that the Borel reinterpretations of A and $\omega^\omega \setminus A$ will be complements in any forcing extension by \mathbb{P}_{α_*} . We have to show that if

- p is a condition in \mathbb{P}_{α_*} ,
- $A \subseteq \omega^\omega$ is universally measurable and
- τ is a \mathbb{P}_{α_*} -name for a Borel measure on ω^ω

then there exist a condition $p' \leq p$ and a \mathbb{P}_{α_*} -name \dot{B} for a Borel subset of ω^ω such that p' forces the symmetric difference of \dot{B} and the Borel reinterpretation of A to be τ -null. By 211X(e) of [3] it suffices to consider the case where τ is a name for a probability measure. We let $P(\omega^\omega)$ be the space of Borel probability measures on ω^ω , with the topology given in Section 17.E of [7].

Fix p , A and τ as in the previous paragraph, and fix in addition X , \bar{Y} , \bar{i} and \bar{D} (with $\tau \in X$) such that $S = \langle \bar{C}, F, \mathbf{q}, p, X, \bar{Y}, \bar{i}, \bar{D} \rangle$ is a suitable data sequence. By Lemma 3.5, we may fix in addition a condition tree

$$(T, \langle p_\rho : \rho \in T \rangle)$$

relative to S . There is then a continuous function f from $[T]$ to $P(\omega^\omega)$ such that each value $f(x)$ is the realization of τ by $\{p_x \upharpoonright_n : n \in \omega\}$. Let λ be Lebesgue measure for $[T]$ and let ν be the measure on ω^ω defined by setting $\nu(B)$ to be $\int_{[T]} f(x)(B) d\lambda$. Since A is universally measurable, there exist Borel sets B and E contained in ω^ω such that E is ν -null and $A \triangle B \subseteq E$. The same relationship will then hold for the (Borel) reinterpretations of A , B and E in any forcing extension by \mathbb{P}_{α_*} . Let K be a closed non-null set of $x \in [T]$ for which E is $f(x)$ -null.

The statement that E is $f(x)$ -null for each $x \in K$ is Π_2^1 in codes for f , E and K , and therefore absolute. One way to see this is to note that for any wellfounded model M of a sufficiently large fragment of ZFC containing these codes and some $x \in K$, M computes some values for the $f(x)$ -measures of E and its complement, and it must compute them correctly.

We can then apply Lemma 5.4 for K in the case $i = 0$, $j = i_*$. The resulting condition q' then forces that the realization of τ in the \mathbb{P}_{α_*} -extension will be in the f -image of K . Since the reinterpretation of E will be $f(x)$ -null in this extension for all x in the reinterpretation of K , the reinterpretations of B and E will witness that the Borel reinterpretation of A is measurable relative to the realization of τ . \square

Essentially the same argument shows that extensions by partial orders of the form \mathbb{P}_{α_*} are measured, in the sense of [10].

6.2 Remark. A set $A \subseteq \omega^\omega$ is universally null if $f^{-1}[A]$ is Lebesgue-null whenever $f: \omega^\omega \rightarrow \omega^\omega$ is a Borel function, and universally meager if $f^{-1}[A]$ is meager whenever $f: \omega^\omega \rightarrow \omega^\omega$ is a Borel function. Lemma 5.2 (along with the fact that universally null and universally meager sets cannot contain perfect sets) shows that in the forcing extensions we consider here, all universally null or universally meager sets have cardinality at most \aleph_1 . It is an open question whether the classes of universally null and universally meager sets can coincide. This question is not answered by the arguments in this paper. A standard recursive construction shows that the Continuum Hypothesis (and more generally, Martin's Axiom) implies the existence of nonmeager universally null sets (for instance, Luzin sets, as discussed on page 539 of [6]; see also [9, 1]). Since the iterations considered here preserve nonmeager sets (via Theorem 5.2 of [8] as

cited in Remark 4.4), Theorem 6.1 implies that there exist such sets in the models produced in this paper. Since universally null sets cannot contain perfect sets, a nonmeager universally null set cannot have the property of Baire. This shows that there are universally measurable sets in the extensions produced in this paper which are not universally categorical.

6.3 Remark. When $a \subseteq \omega$, the inner model $\mathbf{L}[a]$ contains a $\Delta_2^1(a)$ wellordering of its reals which remains $\Delta_2^1(a)$ in any outer model. Since the iterations considered here preserve non-null sets (via Theorem 6.3 and Lemma 6.4 of [8] as cited in Remark 4.4), the reals of $\mathbf{L}[a]$ form a non-null set in the models considered here. By [13], any set of reals wellordered by a universally measurable relation is universally null. It follows in the extensions produced in this paper, the canonical wellordering of the reals constructed in $\mathbf{L}[a]$ is a $\Delta_2^1(a)$ set which is not universally measurable. The analogous results for category ([8] and [13] again) show that this set is also not universally categorical.

References

- [1] J. Brendle, P. Larson, *Nonmeager universally null sets*, preprint, 2017
- [2] Q. Feng, Qi, M. Magidor, H. Woodin, *Universally Baire sets of reals*. In : **Set theory of the continuum**, Math. Sci. Res. Inst. Publ., 26, Springer, New York, 1992, 203–242
- [3] D.H. Fremlin, **Measure Theory**, Volume Two, Torres Fremlin, Colchester, 2003
- [4] D.H. Fremlin, **Measure Theory**, Volume Four, Part I, Torres Fremlin, Colchester, 2006
- [5] D.H. Fremlin, *Problems*, available at <https://www1.essex.ac.uk/maths/people/fremlin/problems.pdf>
- [6] T. Jech, **Set Theory**, Springer, 2003
- [7] A. Kechris, **Classical Descriptive Set Theory**, Springer-Verlag, 1995
- [8] J. Kellner, S. Shelah, *Preserving preservation*, J. Symbolic Logic 70 (2005) 3, 914–945
- [9] P.B. Larson, I. Neeman, S. Shelah, *Universally measurable sets in generic extensions*, Fund. Math. 208 (2010) 2, 173–192
- [10] P.B. Larson, J. Zapletal, *Polar forcings and measured extensions*, Topology Appl. 323 (2023), Paper No. 108290
- [11] T. Nishiura, **Absolute Measurable Spaces**, Encyclopedia of Mathematics and its Applications, Cambridge University Press, 2008
- [12] J.C. Oxtoby, **Measure and Category**, Springer-Verlag, 1980

- [13] I. Reclaw, *On a construction of universally null sets*, Real Anal. Exchange 27 (2002) 1, 321–324
- [14] S. Shelah, *Whitehead groups may be not free, even assuming CH, I* Israel J. Math 28 (1977) 3, 193–204
- [15] S. Shelah, *Whitehead groups may be not free, even assuming CH, II* Israel J. Math 35 (1980) 3, 257–285
- [16] S. Shelah, **Proper and Improper Forcing**, Springer, 1998
- [17] S. Shelah, W.H. Woodin, *Large cardinals imply that every reasonably definable set of reals is Lebesgue measurable*, Israel J. Math. 70 (1990) 3, 381–394